

# ESC195 Midterm 2 Review

QiLin Xue

March 31, 2021

## Contents

<b>1 Sequences and Series</b>	<b>1</b>
1.1 Important Limits	2
1.2 Series	2
1.3 Convergence Tests	3
<b>2 Power Series</b>	<b>4</b>
2.1 Binomial Theorem	5
2.2 Important Series	5
<b>3 Fourier Series</b>	<b>6</b>
<b>4 Vector Functions</b>	<b>6</b>
4.1 Curvature	6
<b>5 Limits of Multivariable Functions</b>	<b>7</b>
5.1 Delta-Epsilon Proofs	7
<b>6 Partial Derivatives</b>	<b>8</b>
<b>7 Other Important Things</b>	<b>8</b>

## 1 Sequences and Series

A sequence  $\{a_n\}$  is:

- increasing iff  $a_n < a_{n+1}$
- non-decreasing iff  $a_k \leq a_{n+1}$
- decreasing iff  $a_n > a_{n+1}$
- non-increasing iff  $a_n \geq a_{n+1}$

**Definition:** We can define  $\lim_{n \rightarrow \infty} a_n = L$  iff for every  $\epsilon > 0$ , there exists an integer  $k > 0$  such that if  $n \geq k$ , then  $|a_n - L| < \epsilon$ .

**Example 1:** Let us prove  $\lim_{n \rightarrow \infty} \frac{n}{n+1} = 1$ . We find  $k$  such that  $\left| \frac{n}{n+1} - 1 \right| < \epsilon$  for  $n \geq k$ . This can be rewritten as:

$$\left| \frac{1}{n+1} \right| < \epsilon \quad (1)$$

or  $|n+1| > \frac{1}{\epsilon}$ . Thus, if we choose  $k = \frac{1}{\epsilon}$  such that if we choose  $n > k = \frac{1}{\epsilon}$ , then:

$$\left| \frac{n}{n+1} - 1 \right| = \left| \frac{1}{n+1} \right| < \left| \frac{1}{n} \right| < \frac{1}{k} = \epsilon \quad (2)$$

Therefore,  $\lim_{n \rightarrow \infty} \frac{n}{n+1} = 1$ .

**Theorem: Uniqueness of a Limit:** If  $\lim_{n \rightarrow \infty} a_n = L$  and  $\lim_{n \rightarrow \infty} a_n = M$ , then  $L = M$ .

**Theorem: Monotonic Sequence Theorem:** A bounded nondecreasing sequence converges to its least upper bound. A bounded non increasing sequence converges to its greatest lower bound.

**Theorem: Pinching Theorem for Sequences:** If for large  $n$ ,  $a_n \leq b_n \leq c_n$  and if  $\lim_{n \rightarrow \infty} a_n = L$  and  $\lim_{n \rightarrow \infty} c_n = L$ , then  $\lim_{n \rightarrow \infty} b_n = L$ .

If a sequence has a limit, it is said to be convergent. Otherwise, it is divergent. This leads to the following:

1. If a sequence is convergent, it is bounded.
2. If a sequence is unbounded, it is divergent.
3. A bounded sequence is not necessarily convergent.

The limit has a few properties. Let  $\lim_{n \rightarrow \infty} a_n = L$  and  $\lim_{n \rightarrow \infty} b_n = M$ . Then:

1.  $\lim_{n \rightarrow \infty} (a_n + b_n) = L + M$
2.  $\lim_{n \rightarrow \infty} \alpha a_n = \alpha L$  for  $\alpha \in \mathbb{R}$ .
3.  $\lim_{n \rightarrow \infty} a_n b_n = L \cdot M$
4.  $\lim_{n \rightarrow \infty} \frac{1}{b_n} = \frac{1}{M}$  for  $b_n \neq 0, M \neq 0$ .
5.  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{L}{M}$  for  $b_n \neq 0, M \neq 0$ .

## 1.1 Important Limits

- For  $x > 0$ ,  $\lim_{n \rightarrow \infty} x^{1/n} = 1$ .
- If  $|x| < 1$ , then  $\lim_{n \rightarrow \infty} x^n = 0$ .
- For  $\alpha > 0$ ,  $\lim_{n \rightarrow \infty} \frac{1}{n^\alpha} = 0$ .
- $\lim_{n \rightarrow \infty} \frac{x^n}{n!} = 0$  for  $x \in \mathbb{R}$ .
- $\lim_{n \rightarrow \infty} \frac{n!}{n^n} = 0$
- $\lim_{n \rightarrow \infty} \frac{\ln n}{n} = 0$ .
- $\lim_{n \rightarrow \infty} n^{1/n} = 1$ .
- $\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x$

## 1.2 Series

You can expect the limit laws to be familiar:

- If  $\sum_{k=0}^{\infty} a_k = L$  and  $\sum_{k=0}^{\infty} b_k = M$ , then  $\sum_{k=0}^{\infty} (a_k + b_k) = L + M$ .

- If  $\sum_{k=0}^{\infty} a_k = L$ , then  $\sum_{k=0}^{\infty} \alpha a_k = \alpha L$  for  $\alpha \in \mathbb{R}$ .

Oftentimes, we wish to set the lower bound to a higher number to do a proof (i.e. to bound a function). Then:

**Theorem:** If  $\sum_{k=0}^{\infty} a_k$  converges iff  $\sum_{k=j}^{\infty} a_k$  converges where  $j$  is a positive integer.

**Theorem:** If  $\sum_{k=0}^{\infty} a_k$  converges, then  $a_k \rightarrow 0$  as  $k \rightarrow \infty$ . Taking the contraposition, we have that if  $a_k \not\rightarrow 0$  as  $k \rightarrow \infty$ , then  $\sum_{k=0}^{\infty} a_k$  diverges.

Note that the inverse isn't necessarily true. If  $a_k \rightarrow 0$  as  $k \rightarrow \infty$ , the sum does not necessarily converge.

### 1.3 Convergence Tests

Here are the following convergence tests:

**Theorem: Integral Test:** If  $f$  is continuous, decreasing, and positive on  $[1, \infty)$ , then:  $\sum_{k=1}^{\infty} f(k)$  converges if and only if  $\int_1^{\infty} f(x) dx$  converges.

**Theorem: P Series:** The  $p$ -series is:

$$\sum_{k=1}^{\infty} \frac{1}{k^p} \quad (3)$$

which will converge if  $p > 1$  since  $\int_1^{\infty} \frac{dx}{x^p}$  converges iff  $p > 1$ .

**Theorem: Direct Comparison Test:** Given  $\sum a_k$  and  $\sum b_k$  with  $a_k > 0$  and  $b_k > 0$ :

1. If  $\sum b_k$  is convergent, and if  $a_k \leq b_k$  for all sufficiently large  $k$ , then  $\sum a_k$  converges.
2. If  $\sum b_k$  is divergent and  $a_k > b_k$  for all  $k$  sufficiently large, then  $\sum a_k$  diverges.

**Theorem: Limit Comparison Test:** Given  $\sum a_k$ ,  $\sum b_k$  where  $a_k > 0$  and  $b_k > 0$ :

1. If  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c > 0$ , then both series converge or diverge.
2. If  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$  and if  $\sum b_n$  converges, then  $\sum a_n$  converges.
3. If  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \infty$  and if  $\sum b_n$  diverges, then  $\sum a_n$  diverges.

**Theorem: Alternating Series Test:** Let  $\{a_k\}$  be a sequence of positive numbers. If and only if  $a_{k+1} < a_k$  and  $a_k \rightarrow 0$  as  $k \rightarrow \infty$ , then:

$$\sum_{k=1}^{\infty} (-1)^{k-1} a_k \quad (4)$$

converges.

**Theorem: Absolute Convergence Test:** If  $\sum |a_k|$  converges, then  $\sum a_k$  converges.

If  $\sum |a_k|$  converges, we say that  $\sum a_k$  is absolutely convergent. If  $\sum a_k$  converges, but  $\sum |a_k|$  does not, we say  $\sum a_k$  is conditionally convergent.

**Theorem: Root Test:** Given  $\sum a_k$ ,  $a_k \geq 0$ . If  $(a_k)^{1/k} \rightarrow p$  as  $k \rightarrow \infty$ , then:

1. If  $p < 1$ , then  $\sum a_k$  converges.
2. If  $p > 1$ , then  $\sum a_k$  diverges.
3. If  $p = 1$  the test is inconclusive.

**Theorem: Ratio test:** Given  $\sum a_k$ , with  $a_k > 0$ . If  $\frac{a_{k+1}}{a_k} \rightarrow \lambda$  as  $k \rightarrow \infty$ , then:

1. If  $\lambda < 1$ ,  $\sum a_k$  converges.
2. If  $\lambda > 1$ ,  $\sum a_k$  diverges.
3. If  $\lambda = 1$ , the test is inconclusive.

If in doubt, follow this (incomplete) check-list:

1. Check if the *sequence* diverges.
2. If it's a power series, use a ratio test.
3. Use comparison + p-series test to bound a sequence by  $\frac{1}{n^p}$ .
4. Factor?

## 2 Power Series

A power series is a series in the form:

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \dots \quad (5)$$

The following gives the Taylor series. If  $a = 0$ , we have a special case, known as the Maclaurin series.

**Theorem:** If  $f(x)$  has a power series representation about  $a$ :

$$f(x) = \sum_{n=0}^{\infty} c_n (x - a)^n \quad (6)$$

with  $|x - a| < R$ . Then the coefficients of the series are  $c_n = \frac{f^{(n)}(a)}{n!}$ .

The  $n$ th degree Taylor polynomial of  $f$  about  $a$  can be written as:

$$T_n(x) = \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (x - a)^i = f(a) + \frac{f'(a)}{1!} (x - a) + \dots + \frac{f^{(n)}(a)}{n!} (x - a)^n \quad (7)$$

This comes with some remainder, which can be calculated below:

**Theorem:** If  $f(x) = T_n(x) + R_n(x)$  and  $\lim_{n \rightarrow \infty} R_n(x) = 0$  for  $|x - a| < R$ . Then  $f$  is equal to the sum of its Taylor series.

Given that  $f$  has  $n + 1$  continuous derivatives on an open interval  $I$  containing  $a$ , then for all  $x \in I$ :

$$f(x) = f(a) + f'(a)(x - a) + \frac{f''(a)(x - a)^2}{2!} + \cdots + \frac{f^{(n)}(a)(x - a)^n}{n!} + R_n(x) \quad (8)$$

where

$$R_n(x) = \frac{1}{n!} \int_a^x f^{(n+1)}(t)(x - t)^n dt \quad (9)$$

This is often a difficult integral to calculate, but we can set the upper bound using the mean value theorem:

$$|R_n| \leq \frac{f^{(n+1)}(z)|x - a|^{n+1}}{(n + 1)!} \quad (10)$$

where  $a \leq z \leq x$  is chosen to maximize  $f^{(n+1)}(z)$ . For an alternating series, we have:

$$R_n < |a_{n+1}| \quad (11)$$

## 2.1 Binomial Theorem

The binomial theorem tells us:

$$(a + b)^k = a^k + ka^{k-1}b + \frac{k(k-1)}{2!}a^{k-2}b^2 + \cdots + \frac{k(k-1)(k-2)\cdots(k-n+1)}{k!}a^{k-n}b^n \quad (12)$$

$$= \sum_{n=0}^k \binom{k}{n} a^{k-n} b^n \quad (13)$$

## 2.2 Important Series

- $e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{3} + \cdots = \sum_{n=0}^{\infty} \frac{x^n}{n!}$  where  $I = (-\infty, \infty)$
- $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^6}{7!} + \cdots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$  where  $I = (-\infty, \infty)$ .
- $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$  where  $I = (-\infty, \infty)$ .
- $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n}$  for  $I = (-2, 1]$ .
- $\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots = \sum_{n=0}^{\infty} x^n$  for  $I = (-1, 1)$ .
- $\tan^{-1}(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$  for  $[-1, 1]$ .

### 3 Fourier Series

The big idea is to write a periodic function in terms of a trigonometric basis:

**Theorem:** For  $f(t)$  periodic, with fundamental period  $T$ , continuous and piecewise differentiable, then:

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos(n\omega t) + b_n \sin(n\omega t) \quad (14)$$

where  $\omega = \frac{2\pi}{T}$  is known as the Fourier series of  $f$ .  $a_n$  and  $b_n$  are Fourier coefficients. The coefficients are given by:

$$a_n = \frac{2}{T} \int_{-T/2}^{T/2} f(t) \cos(n\omega t) dt \quad (15)$$

$$b_n = \frac{2}{T} \int_{-T/2}^{T/2} f(t) \sin(n\omega t) dt \quad (16)$$

for  $n = 1, 2, 3, \dots$

You can apply the following shortcuts:

- If  $f(t)$  is odd, then  $a_n = 0$ .
- If  $f(t)$  is even, then  $b_n = 0$ .

### 4 Vector Functions

For a vector function  $\vec{f}(t)$  which maps a scalar to a vector, the operate and differentiate the way you'll expect. We have the following differentiation rules:

- $(\vec{f} + \vec{g})'(t) = \vec{f}'(t) + \vec{g}'(t)$
- $(\alpha \vec{f})'(t) = \alpha f'(t)$
- $(u\vec{f})'(t) = u(t)\vec{f}'(t) + u'(t)\vec{f}(t)$
- $(\vec{f} \cdot \vec{g})'(t) = [\vec{f}(t) \cdot \vec{g}'(t)] + [\vec{f}'(t) \cdot \vec{g}(t)]$
- $(\vec{f} \times \vec{g})'(t) = [\vec{f}'(t) \times \vec{g}(t)] + [\vec{f}(t) \times \vec{g}'(t)]$
- $(\vec{f} \circ u)'(t) = \vec{f}'(u(t))u'(t)$

**Definition:** Let  $C$  be parametrized by  $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$  and be differentiable. Then  $\vec{r}'(t) = x'(t)\hat{i} + y'(t)\hat{j} + z'(t)\hat{k}$  if not  $\vec{0}$ , is tangent to the curve  $C$  at the point  $P(x(t), y(t), z(t))$  and  $\vec{r}'(t)$  points in the direction of increasing  $t$ .

The arclength is:

$$s = \int_a^b \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2} dt \quad (17)$$

#### 4.1 Curvature

In two dimensions, the curvature of a *2-dimensional curve* is defined as:

$$\kappa = \left| \frac{d\phi}{ds} \right| \quad (18)$$

where:

$$\frac{dy}{dx} = y' = \tan \phi \implies \phi = \tan^{-1}(y') \quad (19)$$

and the radius of curvature is:

$$r = \frac{1}{\kappa}. \quad (20)$$

In three dimensions, there are three ways of calculating curvature:

- Let  $\vec{T}$  be the unit tangent  $\vec{T} = \frac{\vec{r}'(t)}{\|\vec{r}'(t)\|}$  that points in the direction of the curve.

$$\kappa = \left\| \frac{d\vec{T}}{ds} \right\| \quad (21)$$

- For a curve  $\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$ , we have:

$$\kappa = \left\| \frac{d\vec{T}}{dt} \cdot \frac{dt}{ds} \right\| = \frac{\|\vec{T}'\|}{\|\vec{r}'\|} \quad (22)$$

- We can also define it using the cross product:

$$\kappa = \frac{\|\vec{r}''(t) \times \vec{r}'(t)\|}{\|\vec{r}'(t)\|^3} \quad (23)$$

The **binormal vector** is given by:

$$\vec{B}(t) = \vec{T} \times \vec{N} \quad (24)$$

and gives the vector normal to the osculating plane. Here, the normal vector is:

$$\vec{N}(t) = \frac{\vec{T}'(t)}{\|\vec{T}'\|} \quad (25)$$

## 5 Limits of Multivariable Functions

Suppose we have the limit:

$$\lim_{(x,y) \rightarrow (0,0)} \frac{f(x,y)}{g(x,y)} \quad (26)$$

where  $f(0,0) = g(0,0) = 0$ . Here are a few strategies to show if it exists or not:

- If in doubt, test if it doesn't exist first. Try the path  $x = 0$ ,  $y = 0$ , and  $y = mx$ .
- The limit will most likely not exist if the order of the denominator is higher than the numerator, and it will likely exist if the order of the numerator is higher than the denominator.
- Use the squeeze theorem. One helpful strategy is to have the upper and lower bound be in terms of one variable.

### 5.1 Delta-Epsilon Proofs

Helpful tips:

- Use triangle inequality.
- Write  $\delta$  restriction as a scalar function (i.e. square root)
- Attempt to bound functions of two variables in terms of a single variable.

**Example 2:** Suppose we wish to prove that:

$$\lim_{(x,y) \rightarrow (1,1)} (x + y) = 2 \quad (27)$$

For any  $\epsilon > 0$  such that  $|f(x,y) - f(1,1)| < \epsilon$ , we can pick a  $\delta > 0$  where  $\|(x,y) - (1,1)\| < \delta$ . We can write this last statement as:

$$\sqrt{(x-1)^2 + (y-1)^2} < \delta \quad (28)$$

Using the triangle inequality, we can write:

$$|f(x, y) - f(1, 1)| = |x + y - 2| \quad (29)$$

$$= |(x - 1) + (y - 1)| \quad (30)$$

$$\leq |x - 1| + |y - 1| \quad (31)$$

$$= \sqrt{(x - 1)^2} + \sqrt{(y - 1)^2} \quad (32)$$

$$\leq \sqrt{(x - 1)^2 + (y - 1)^2} + \sqrt{(y - 1)^2 + (x - 1)^2} \quad (33)$$

$$= 2\sqrt{(x - 1)^2 + (y - 1)^2} \quad (34)$$

$$< 2\delta = \epsilon \quad (35)$$

where we used our statement from earlier for the last line. Therefore, we need to pick  $\delta = \frac{\epsilon}{2}$ .

## 6 Partial Derivatives

The partial derivative of  $f(x, y)$  is given by:

$$f_x(x, y) = \frac{\partial}{\partial x} f(x, y) = \lim_{h \rightarrow 0} \frac{f(x + h, y) - f(x, y)}{h} \quad (36)$$

or for the partial derivative with respect to  $y$ :

$$f_y(x, y) = \frac{\partial}{\partial y} f(x, y) = \lim_{h \rightarrow 0} \frac{f(x, y + h) - f(x, y)}{h} \quad (37)$$

This can be extended to an arbitrary number of dimensions. We can also have mixed partials, such as:

$$\frac{\partial}{\partial x} \frac{\partial f}{\partial x} \rightarrow \frac{\partial^2 f}{\partial x^2} \quad (38)$$

$$\frac{\partial}{\partial y} \frac{\partial f}{\partial x} \rightarrow \frac{\partial^2 f}{\partial y \partial x} \quad (39)$$

**Theorem:** Clairaut's Theorem says that:

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial^2 f}{\partial x \partial y} \quad (40)$$

on every open set on which  $f$  and its partials  $\frac{\partial f}{\partial x}$ ,  $\frac{\partial f}{\partial y}$ ,  $\frac{\partial^2 f}{\partial x \partial y}$ ,  $\frac{\partial^2 f}{\partial y \partial x}$  are continuous.

## 7 Other Important Things

- $\cos(n\pi) = (-1)^n$
- $\cos\left(\frac{n\pi}{2}\right) = \begin{cases} (-1)^{n/2} & n \text{ is even} \\ 0 & n \text{ is odd} \end{cases}$
- $\sin\left(\frac{n\pi}{2}\right) = \begin{cases} 0 & n \text{ is even} \\ (-1)^{\frac{n-1}{2}} & n \text{ is odd} \end{cases}$
- $(-1)^n + (-1)^{n+1} = 0$
- $1 + (-1)^n = \begin{cases} 2 & \text{even} \\ 0 & \text{odd} \end{cases}$
- $(-1)^{-n} = (-1)^n$
- $(-1)^{1-n} = (-1)^{n+1}$