

# PHY294: Review Sheet

QiLin Xue

Winter 2022

## 1 The Three-Dimensional Schrödinger Equation.

In general, the Schrödinger Equation can be written as

$$\nabla^2 \psi = \frac{2M}{\hbar^2} [U - E] \psi \quad (1)$$

where  $\nabla^2$  is the Laplacian, which is a fancy way of writing:

$$\nabla^2 \psi = \frac{\partial^2 \psi}{\partial x^2} + \frac{\partial^2 \psi}{\partial y^2} + \frac{\partial^2 \psi}{\partial z^2} \quad (\text{Cartesian})$$

$$= \frac{\partial^2 \psi}{\partial r^2} + \frac{1}{r} \frac{\partial \psi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \psi}{\partial \phi^2} + \frac{\partial^2 \psi}{\partial z^2} \quad (\text{Cylindrical})$$

$$= \frac{1}{r} \frac{\partial^2}{\partial r^2} (r\psi) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial \psi}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \psi}{\partial \phi^2}$$

### 1.1 Particle in a Box

For a particle in an infinite square well, the allowed energies are

$$E = \frac{\hbar^2 \pi^2}{2M} \left( \frac{n_x^2}{L_x^2} + \frac{n_y^2}{L_y^2} + \frac{n_z^2}{L_z^2} \right) \quad (2)$$

Here,  $n_x, n_y, n_z$  are known as **quantum numbers**. When the same energy occurs at different quantum numbers, then we call it a **degeneracy**.

### 1.2 Central Force Problem

#### 2-Dimensions

For the two-dimensional case, our solution is in the form of  $\psi(r, \theta) = R(r)\Theta(\theta)$ . Solving for  $\Theta(\theta)$  gives:

$$\Phi(\phi) = A \sin(m\phi) + B \cos(m\phi),$$

which must satisfy the property  $\Phi(\phi) = \Phi(\phi + 2\pi)$ , forcing  $m$  to be an integer. The radial part  $R(r)$  cannot be solved since it depends on the specific potential.

#### 3-Dimensions

The solution is in the form of  $\phi(r, \theta, \phi) = R(r)\Theta(\theta)\Phi(\phi)$ .

- Solving for  $\Phi(\phi)$  gives the same equation as the  $\Theta(\theta)$  case in 2-D, leading to the same quantum numbers.
- Solving for  $\Theta(\theta)$  gives a parameter  $k = l(l+1)$  where  $l \geq |m|$  and is an integer.

### 1.3 Angular Momentum

The magnitude of the angular momentum is quantized and given by

$$L = \sqrt{l(l+1)}\hbar. \quad (3)$$

The angular momentum acts in a certain direction away from the  $z$ -axis, and

$$L_z = m\hbar, \quad (4)$$

where  $m$  has the usual restrictions. As a result, we can often draw these as vectors on a diagram. For a given  $l$ , there are  $2l + 1$  possible vectors.

## 1.4 Hydrogen Atom

Consider the simplest (yet useful) potential  $U(r) = \frac{-ke^2}{r}$ , for a hydrogen atom. The  $R(r)$  part is given by

$$\frac{d^2}{dr^2}(rR) = \frac{2m_e}{\hbar^2} \left[ -\frac{ke^2}{r} + \frac{l(l+1)\hbar^2}{2m_e r^2} - E \right] (rR), \quad (5)$$

which only has acceptable solutions if

$$E = -\frac{m_e(ke^2)^2}{2\hbar^2} \frac{1}{n^2} = -\frac{E_R}{n^2}, \quad (6)$$

where  $n$  has the restriction  $n > l$ .

## 1.5 Hydrogenic Wave Functions

The solution for  $R(r)$  is in the form of

$$R(r) = Ae^{-r/(na_B)} \quad (7)$$

where  $a_B = \frac{\hbar^2}{m_e ke^2}$  is the Bohr radius, and  $A$  is some function. The radial probability density is

$$P(r) = 4\pi r^2 |R(r)|^2 \quad (8)$$

## 1.6 Hydrogen-like Ions

We can extend this discussion to any ion with one electron bounded to a nucleus of charge  $Ze$ . Note that  $\Phi, \Theta$  are left unchanged and everything else can be derived by replacing  $ke^2$  with  $Zke^2$  and as a direct consequence, replacing  $a_B$  with  $a_B/Z$ .